

# McMaster University

## Advanced Optimization Laboratory



### **Title:**

On a conjecture of Erdős for multiplicities of cliques

### **Authors:**

Antoine Deza, Frantisek Franek, and Min Jing Liu

ADVANCED OPTIMIZATION LABORATORY,  
DEPARTMENT OF COMPUTING AND SOFTWARE,  
MCMASTER UNIVERSITY, HAMILTON, ONTARIO, CANADA.  
*Email:* deza, franek, liumj@mcmaster.ca

**AdvOL-Report No. 2010/4**

May 2010, Hamilton, Ontario, Canada

# On a conjecture of Erdős for multiplicities of cliques

Antoine Deza, Frantisek Franek, and Min Jing Liu \*

Advanced Optimization Laboratory  
Department of Computing and Software  
McMaster University, Hamilton, Ontario, Canada L8S 4K1  
email: {deza,franek,liumj}@mcmaster.ca

**Abstract.** Denote by  $k_t(G)$  the number of cliques of order  $t$  in a graph  $G$  having  $n$  vertices. Let  $k_t(n) = \min\{k_t(G) + k_t(\overline{G})\}$  where  $\overline{G}$  denotes the complement of  $G$ . Let  $c_t(n) = k_t(n)/\binom{n}{t}$  and  $c_t$  be the limit of  $c_t(n)$  for  $n$  going to infinity. A 1962 conjecture of Erdős stating that  $c_t = 2^{1-\binom{t}{2}}$  was disproved by Thomason in 1989 for all  $t \geq 4$ . Tighter counterexamples have been constructed by Jagger, Šťovíček and Thomason in 1996, by Thomason for  $t \leq 6$  in 1997, and by Franek for  $t = 6$  in 2002. We investigate a computational framework to search for tighter upper bounds for small  $t$  and provide the following improved upper bounds for  $t = 6, 7$  and  $8$ :  $c_6 \leq 0.7445 \times 2^{1-\binom{6}{2}}$ ,  $c_7 \leq 0.6869 \times 2^{1-\binom{7}{2}}$ , and  $c_8 \leq 0.7002 \times 2^{1-\binom{8}{2}}$ . The constructions are based on a large but highly regular variant of Cayley graphs for which the number of cliques and cocliques can be expressed in closed form. Note that if we consider the quantity  $e_t = 2^{\binom{t}{2}-1}c_t$ , the new upper bound of 0.687 for  $e_7$  is the first bound for any  $e_t$  smaller than the lower bound of 0.695 for  $e_4$  due to Giraud in 1979.

**Keywords:** clique, coclique, Cayley graph, complete graph, subgraph

## 1 Introduction

Denote by  $k_t(G)$  the number of cliques of order  $t$  in a graph  $G$  having  $n$  vertices. Let  $k_t(n) = \min\{k_t(G) + k_t(\overline{G})\}$  where  $\overline{G}$  denotes the complement of  $G$ . The cliques in  $\overline{G}$  are referred to as cocliques. If we want to be specific about their sizes, we talk of  $t$ -cliques and  $t$ -cocliques. Let  $c_t(n) = k_t(n)/\binom{n}{t}$  and  $c_t = \lim_{n \rightarrow \infty} c_t(n)$ . Since we can view  $G$  and  $\overline{G}$  as a 2-colouring of the edges of the complete graph  $K_n$ ,  $c_t(n)$  denotes the minimum proportion of monochromatic  $t$ -cliques and  $t$ -cocliques for all 2-colourings of the edges of  $K_n$ .

A conjecture of Erdős related to Ramsey's Theorem [2], states that  $c_t = 2^{1-\binom{t}{2}}$ . The conjecture is clearly true for  $t = 2$ , and using Goodman's approach [8], one can show that the conjecture holds for  $t = 3$ . One of the motivations behind the conjecture is the fact that the conjecture holds for any  $t$  for random graphs. Erdős and Moon [3] showed that a modified conjecture for complete bipartite subgraphs of bipartite graphs is true. Sidorenko [11] showed that a modified conjecture for cycles is true, but not true for certain incomplete subgraphs. Franek and Rödl [5] showed that the original conjecture for  $t = 4$  is true for nearly quasirandom, and hence quasirandom graphs.

---

\* corresponding author

Thomason [12] disproved the conjecture for  $t \geq 4$  by exhibiting constructions achieving low numbers of cliques and cocliques. Thomason's upper bounds from [12] were:  $c_4 \leq 0.976 \times 2^{1-\binom{4}{2}}$ ,  $c_5 \leq 0.906 \times 2^{1-\binom{5}{2}}$ , and  $c_t \leq 0.936 \times 2^{1-\binom{t}{2}}$  for  $t \geq 6$ . These bounds were further improved in [13] to  $c_4 \leq 0.9693 \times 2^{1-\binom{4}{2}}$  and  $c_5 \leq 0.8801 \times 2^{1-\binom{5}{2}}$ , in [4] to  $c_6 \leq 0.7446 \times 2^{1-\binom{6}{2}}$ , and in [9] to  $c_t \leq 0.835 \times 2^{1-\binom{t}{2}}$  for  $t \geq 7$ . The construction used in [4] to bound  $c_6$  is based on the approach used by Franek and Rödl [6], who tied the best upper bound for  $c_4$ . It improves the best upper bound for  $t = 7$  to  $c_7 \leq 0.7156 \times 2^{1-\binom{7}{2}}$ . This bound for  $c_7$  was mentioned in a referee report but never formally put forward.

In this paper we investigate a computational framework to search for tighter upper bounds for small  $t$  and give improved upper bounds for  $t = 6, 7$  and  $8$ :  $c_6 \leq 0.74444 \times 2^{1-\binom{6}{2}}$ ,  $c_7 \leq 0.6869 \times 2^{1-\binom{7}{2}}$ , and  $c_8 \leq 0.7002 \times 2^{1-\binom{8}{2}}$ .

Concerning the lower bound, see Conlon [1] for a recent improvement over Erdős's original application of Ramsey's Theorem, and Giraud [7] who showed that  $c_4 \geq 0.695 \times 2^{1-\binom{4}{2}}$ . Note that if we consider the quantity  $e_t = 2^{\binom{t}{2}-1} c_t$ , the new upper bound for  $e_7$  is in fact smaller than Giraud's lower bound for  $e_4$ ; this is the first such upper bound for any  $e_t$ .

## 2 Constructing Counterexamples

In order to improve the upper bound for  $c_t$  for small  $t$ , we follow the approach used in [4, 6] and work with graphs for which the number of cliques and cocliques can be expressed in closed form. This allows a viable search among them for the ones that exhibit the lowest numbers of cliques and cocliques.

For a positive integer  $X$  and  $F \subseteq \{1, 2, \dots, X\}$ , we consider the graph  $G_{X,F}$  whose vertices are all  $2^X$  subsets of  $\{0, 1, \dots, X-1\}$ , and two subsets  $x_i$  and  $x_j$  of  $\{0, 1, \dots, X-1\}$  are connected by an edge if and only if  $|x_i \Delta x_j| \in F$ , where  $\Delta$  denotes the operation of symmetric difference. We clearly have  $\overline{G}_{X,F} = G_{X,\overline{F}}$  where  $\overline{F} = \{1, 2, \dots, X\} - F$ .

Since it would be too complicated to count cliques in  $G_{X,F}$ , we introduce the notion of  $(X, F)$ -tuples and count the  $(X, F)$ -tuples instead. Lemma 1 recalls the straightforward relationship between these quantities. For  $m \geq 1$ , an ordered  $m$ -tuple  $\langle x_0, x_1, \dots, x_{m-1} \rangle$  is an  $(X, F)$ - $m$ -tuple if  $x_i \subseteq \{0, 1, \dots, X-1\}$ ,  $|x_i| \in F$  for  $i < m$ , and  $|x_i \Delta x_j| \in F$  for all  $i \neq j < m$ .

**Lemma 1.** *The number  $k_{m+1}(G_{X,F})$  of cliques of size  $m+1$  in the graph  $G_{X,F}$  satisfies  $k_{m+1}(G_{X,F}) = \frac{2^n}{(m+1)!} S_m(X, F)$  where  $S_m(X, F)$  is the number of  $(X, F)$ - $m$ -tuples.*

*Proof.* We simply illustrate the cases  $m = 1$  and  $m = 2$ . Case  $m = 1$ : let  $\{x_i, x_j\}$  be a 2-clique; i.e. an edge, in  $G_{X,F}$ . Clearly  $\langle x_i \Delta x_j \rangle$  and  $\langle x_j \Delta x_i \rangle$  are  $(X, F)$ -singletons, so we have 2 distinct  $(X, F)$ -singletons for each 2-clique and  $k_2(G_{X,F}) = \frac{2^n}{2!} S_1(X, F)$ . Case  $m = 2$ : let  $\{x_i, x_j, x_k\}$  be a 3-clique in  $G_{X,F}$ . Clearly  $\langle x_i \Delta x_j, x_i \Delta x_k \rangle$  is an  $(X, F)$ -pair of distinct elements. Considering the permutations of  $i, j, k$  we have 3! distinct  $(X, F)$ -pairs for each 3-clique and  $k_3(G_{X,F}) = \frac{2^n}{3!} S_2(X, F)$ .  $\square$

For a positive integer  $d$  and a graph  $G$  of order  $n$ , the graph  $G^d$  is obtained by replacing each vertex of  $G$  by a  $d$ -clique; therefore  $G^d$  has  $dn$  vertices. Besides the edges within the created  $d$ -cliques, there is an edge between two vertices  $v_i$  and  $v_j$  of  $G^d$  if and only if an edge existed in  $G$  between the two vertices corresponding to the  $d$ -cliques containing  $v_i$  and  $v_j$ ,  $i \neq j$ .

**Lemma 2.** We have  $\lim_{d \rightarrow \infty} \frac{k_7(G^d) + k_7(\overline{G^d})}{\binom{dn}{7}} =$

$$\frac{5040(k_7(G) + k_7(\overline{G})) + 15120k_6(G) + 16800k_5(G) + 8400k_4(G) + 1806k_3(G) + 126k_2(G) + k_1(G)}{n^7}$$

*Proof.* A 7-clique in  $G^d$  may arise from the following seven cases which correspond to the possible partitioning of number 7:  $[\{7\}]$ ,  $[\{1, 6\}\{2, 5\}\{3, 4\}]$ ,  $[\{1, 1, 5\}\{1, 2, 4\}\{1, 3, 3\}\{2, 2, 3\}]$ ,  $[\{1, 1, 1, 4\}\{1, 1, 2, 3\}\{1, 2, 2, 2\}]$ ,  $[\{1, 1, 1, 1, 3\}\{1, 1, 1, 2, 2\}]$ ,  $[\{1, 1, 1, 1, 1, 2\}]$ ,  $[\{1, 1, 1, 1, 1, 1, 1\}]$  – we have grouped them by the number of partitions. For illustration, we count the number  $Q_2(d)$  of cliques corresponding to the decompositions  $[\{1, 6\}\{2, 5\}\{3, 4\}]$ . We have:

$Q_2(d) = \binom{2}{1} \binom{d}{1} \binom{d}{6} + \binom{2}{1} \binom{d}{2} \binom{d}{5} + \binom{2}{1} \binom{d}{3} \binom{d}{4} k_2(G) = (2L_1(d) + 6L_2(d) + 10L_3(d)) \frac{d^7}{6!} k_2(G)$  where  $L_1(d) = (1 - \frac{1}{d})(1 - \frac{2}{d})(1 - \frac{3}{d})(1 - \frac{4}{d})(1 - \frac{5}{d})$ ,  $L_2(d) = (1 - \frac{1}{d})^2(1 - \frac{2}{d})(1 - \frac{3}{d})(1 - \frac{4}{d})$  and  $L_3(d) = (1 - \frac{1}{d})^2(1 - \frac{2}{d})^2(1 - \frac{3}{d})$ . To derive similar formulas for the other partitions is straightforward,

giving  $\lim_{d \rightarrow \infty} \frac{k_7(G^d)}{\binom{dn}{7}} = \frac{5040k_7(G) + 15120k_6(G) + 16800k_5(G) + 8400k_4(G) + 1806k_3(G) + 126k_2(G) + k_1(G)}{n^7}$ .

A 7-coclique can only arise in one way, and thus for the number of 7-cocliques, we get

$$\lim_{d \rightarrow \infty} \frac{k_7(\overline{G^d})}{\binom{dn}{7}} = \frac{5040k_7(\overline{G})}{n^7}. \quad \square$$

*Remark.* In general, the coefficients  $\alpha_{m,t}$  for  $k_m(G)$  in the formula reducing the computation of  $\lim_{d \rightarrow \infty} \frac{k_t(G^d) + k_t(\overline{G^d})}{\binom{dn}{t}}$  to counting cliques and cocliques in the underlying graph  $G$  follow a pattern similar to the Pascal triangle equality as we have  $\alpha_{m,t} = m(\alpha_{m,t-1} + \alpha_{m-1,t-1})$ . See Lemma 2 for the case  $t = 7$ . The coefficients for  $k_m(G)$  and other auxiliary results are available online at [10].

We can set  $G = G_{X,F}$  and then substitute  $k_m(G_{X,F})$  by  $S_{m-1}(X, F)$  using Lemma 1, and restate Lemma 2 as:

**Lemma 3.** For a given pair  $(X, F)$ , we have  $\lim_{d \rightarrow \infty} \frac{k_7(G_{X,F}^d) + k_7(\overline{G_{X,F}^d})}{\binom{d2^n}{7}} =$

$$\frac{S_6(X, F) + S_6(X, \overline{F}) + 21S_5(X, F) + 140S_4(X, F) + 350S_3(X, F) + 301S_2(X, F) + 63S_1(X, F) + 1}{2^{6n-20}}.$$

The approach used in [6] is based on an exhaustive search for a pair  $(X, F)$  achieving a low number of cliques and cocliques for  $t = 4$ . The identified best pair  $(10, \{1, 3, 4, 7, 8, 10\})$  yields a tie for the best upper bound for  $c_4$  and was used to achieve  $c_6 \leq 0.7446 \times 2^{1-\binom{6}{2}}$ . The referee's report for [6] mentioned that the same pair yields  $c_7 \leq 0.7156 \times 2^{1-\binom{7}{2}}$  but this bound was never formally put forward. In this paper we improve the bounds for  $c_t$  for  $t = 6, 7$  and 8.

### 3 Computational Framework

Lemma 3 provides a closed formula for computing a limit of a special sequence of graphs determined by a given pair  $(X, F)$ . If this limit is small enough, it constitutes a counterexample to the conjecture of Erdős. Thus, the computational framework consists of a routine to compute all the required  $S_i(X, F)$ 's for a given pair  $(X, F)$  and a routine performing a search for the best  $(X, F)$ . First, in Section 3.1 we discuss the approach for computing  $S_i(X, F)$  that was used previously in [4, 6]. This approach is rather slow and cannot be employed for  $t > 4$ . That is why only a single pair  $(10, \{1, 3, 4, 7, 8, 10\})$  was used in [4]. Then, in Sections 3.2 and 3.3 we discuss a different approach to the computation of  $S_i(X, F)$ 's referred to as  $m$ -approach, and a further enhancement based on symmetry. These techniques provide a significant speedup allowing an exhaustive search for  $t = 6$  and 7 that was previously intractable.

#### 3.1 Straightforward computation of $S_i$

For simplicity, for a given  $X$ ,  $\hat{X}$  denotes the set  $\{0, 1, \dots, X-1\}$ .

##### Straightforward computation of $S_1(X, F)$

Generate all possible  $x_0 \subseteq \hat{X}$  so that  $|x_0| \in F$ ; then

$$S_1(X, F) = \sum_{|x_0| \in F} \binom{X}{|x_0|}$$

##### Straightforward computation of $S_2(X, F)$

Consider an ordered pair  $\langle x_0, x_1 \rangle$  of mutually distinct subsets of  $\hat{X}$ . Clearly,  $x_0 \cap (x_0 \triangle x_1)$ ,  $x_1 \cap (x_0 \triangle x_1)$  and  $x_0 \cap x_1$  are mutually disjoint. Let  $m_0 = |x_0 \cap (x_0 \triangle x_1)|$ ,  $m_1 = |x_1 \cap (x_0 \triangle x_1)|$  and  $m_{01} = |x_0 \cap x_1|$ . We have  $m_0 + m_{01} = |x_0|$ ,  $m_1 + m_{01} = |x_1|$ , and  $m_0 + m_1 = |x_0 \triangle x_1|$ . In addition, we have  $|x_0|$ ,  $|x_1|$  and  $|x_0 \triangle x_1| \in F$ . Thus, once generating all possible valid solutions  $\langle m_0, m_1, m_{01} \rangle$ , we obtain the value of  $S_2(X, F)$  by:

$$S_2(X, F) = \sum_{\text{all valid } \langle m_0, m_1, m_{01} \rangle} \binom{X}{m_0} \binom{X-m_0}{m_1} \binom{X-m_0-m_1}{m_{01}}$$

##### Straightforward computation of $S_i(X, F)$ for $i > 2$

Similar computations, with increasing computation time, are performed to obtain the values of  $S_i(X, F)$ . We need to consider an ordered  $i$ -tuple  $\langle x_0, x_1, x_2, \dots, x_{i-1} \rangle$  of mutually distinct

subsets of  $\hat{X}$ , and find all the valid solutions  $\langle m_0, m_1, m_2, \dots \rangle$ . Then we can compute the sum of the corresponding binomial coefficients using a dynamically expanded and maintained Pascal triangle. Notice that the total number of the solutions increases rather quickly. In general, we have to consider  $(2^i - 1)$  solutions to compute  $S_i(X, F)$ .

### 3.2 The $m$ -approach to computing $S_i$

In Section 3.1, the  $S_i$  was obtained by finding all valid solutions and computing the sum of the corresponding binomial coefficients, a procedure with an  $O(2^{iX})$  worst-case complexity. Therefore, a more efficient approach is required to speed up the computation.

The following example illustrates how knowing and storing solutions for  $S_{i-1}$ 's can be used to faster obtain solutions for  $S_i$ . For the illustration, we consider computing a solution for  $S_3$  while having  $m^* = \langle m_0^*, m_1^*, m_{01}^* \rangle$  a valid solution for  $S_2$ . We could generate a valid solution  $m = \langle m_0, m_1, m_2, m_{01}, m_{02}, m_{12}, m_{012} \rangle$  for  $S_3$  by reusing  $m^*$ , since  $m_0 + m_{02} = m_0^*$ ,  $m_1 + m_{12} = m_1^*$  and  $m_{01} + m_{012} = m_{01}^*$ . The following constraints can be used to check the validity:  $0 \leq m_0 \leq m_0^*$ ,  $0 \leq m_1 \leq m_1^*$ , and  $0 \leq m_{01} \leq m_{01}^*$ . Recall that  $|x_2|$  should be in  $F$ , and thus we can calculate  $m_2$  directly: as  $m_2 = z - m_{12} - m_{02} - m_{012}$  for some  $z \in F$ . We also need to check the symmetric difference relationships among the  $x_i$ 's. However, we only need to check  $|x_0 \Delta x_2| \in F$  and  $|x_1 \Delta x_2| \in F$ .

*Remark* If  $m^*$  is a valid solution for  $S_i$ , and  $m$  is the corresponding valid solution for  $S_{i+1}$ ,

$$Y^* = \binom{X}{m_0^*} \binom{X-m_0^*}{m_1^*} \binom{X-m_0^*-m_1^*}{m_2^*} \binom{X-m_0^*-m_1^*-m_2^*}{m_3^*} \dots$$

is the corresponding product of the binomial coefficients for  $m^*$ , and

$$Y = \binom{X}{m_0} \binom{X-m_0}{m_1} \binom{X-m_0-m_1}{m_2} \binom{X-m_0-m_1-m_2}{m_3} \dots$$

is the corresponding product of the binomial coefficients for  $m$ , then  $Y = Y^* \binom{m_0^*}{m_0} \binom{m_1^*}{m_1} \dots \binom{m_{01\dots i}^*}{m_{01\dots i}} \binom{X-m_0^*-m_1^*-m_{01}^*-\dots}{m_i}$ .

Similarly, to compute  $S_i$  we only need to consider  $2^{i-1}$   $m$ 's, if we reuse the results from the computation of  $S_{i-1}$ .

### 3.3 Exploiting symmetry

This technique to further speed up the computation of  $S_i$  relies on the inherent symmetries of the  $m_i$ 's. We shall illustrate the technique on  $S_2$ : if  $\langle m_0, m_1, m_{01} \rangle$  is a valid solution for  $S_2$  with  $m_0 \neq m_1$ , then  $\langle m_1, m_0, m_{01} \rangle$  is also a valid solution. Since the products of the corresponding binomial coefficients for those two solutions are the same, we only need to compute the product of the binomial coefficients for one solution and multiply it by 2.

Similarly, the symmetries can be exploited for computing  $S_i$  for  $i \geq 2$ . Thus, one can fix the order of the  $x_i$  and take into account multiplicities by multiplying by the corresponding coefficients. We therefore need, for example for the computation of  $S_7$ , to consider only about 1% of the total number of solutions. Table 1 shows the coefficients used for  $S_4$ . The coefficients for other  $S_i$ 's are available online at [10].

Note that while the determination of  $S_i$  and  $\bar{S}_i$  for the first  $i$ 's is very fast even without exploiting the symmetry, the computational gain increases with  $i$ . Table 2 shows the number of solutions that need to be computed when we used the pair  $(X, F) = (11, \{3, 4, 7, 8, 10, 11\})$  to compute  $S_4, S_5$  and  $S_6$ .

$x_i$ ordering	coefficient
$ x_0  >  x_1  >  x_2  >  x_3 $	$4!$
$ x_0  >  x_1  >  x_2  =  x_3 $	$2 \binom{4}{2}$
$ x_0  >  x_1  =  x_2  >  x_3 $	$2 \binom{4}{2}$
$ x_0  =  x_1  >  x_2  >  x_3 $	$2 \binom{4}{2}$
$ x_0  >  x_1  =  x_2  =  x_3 $	$\binom{4}{3}$
$ x_0  =  x_1  >  x_2  =  x_3 $	$\binom{4}{2}$
$ x_0  =  x_1  =  x_2  >  x_3 $	$\binom{4}{3}$
$ x_0  =  x_1  =  x_2  =  x_3 $	$1$

**Table 1.** Ordering of the  $x_i$ 's and corresponding coefficients for  $S_4$  computation

$i$	# of solutions in $S_i$	# of solutions exploiting symmetry	Ratio	# of solutions in $\bar{S}_i$	# of solutions exploiting symmetry	Ratio
4	15,668	1,813	3.0%	4,477	794	5.9%
5	377,196	17,625	0.5%	86,978	8,214	1.7%
6	9,104,496	160,626	0.08%	1,145,103	55,803	0.46%

**Table 2.** Exploiting symmetry for  $(X, F) = (11, \{3, 4, 7, 8, 10, 11\})$

## 4 New Upper Bounds for $c_6$ , $c_7$ , and $c_8$

Using the approach described in Sections 3.2 and 3.3 we performed an exhaustive search on  $(X, F)$  for  $X = 9, 10, 11$  and  $12$  for  $t = 6$  and  $7$ , using code written in C++.

Besides the usual testing and verification, we also computationally checked the new program by recomputing previously known values as well as theoretically known ones. We first computed the values of  $S_1, \dots, S_6$  for all the previously used pairs  $(X, F)$  and obtained the same results, using a tiny fraction of the computation time previously required. We then computed the values of  $S_1, \dots, S_7$  for *full families* because for such a family  $\{1, 2, \dots, X\}$  the number  $i$ -tuples can be expressed using Lemma 1 with a closed form formula  $S_i = \frac{(2^X - 1)!}{(2^X - i - 1)!}$ . The computed and theoretical values coincided, which is a strong indication that the generation of valid solutions is both sound and complete.

The best results were achieved for  $t = 6$  by  $(X, F) = (10, \{1, 3, 4, 7, 8\})$  yielding  $c_6 \leq 0.74444 \times 2^{1 - \binom{6}{2}}$ , see Table 3. For  $t = 7$  by  $(X, F) = (11, \{3, 4, 7, 8, 10, 11\})$  yielding  $c_7 \leq 0.6869 \times 2^{1 - \binom{7}{2}}$ , see Table 4.

Representing the pair  $(X, F)$  as the characteristic vector of  $F$  as a subset of  $\{1, 2, \dots, X\}$ , one notices the best result for  $t = 6$ , respectively  $t = 7$ , are obtained with  $(X, F) = [1011001100]$ , respectively  $(X, F) = [00110011011]$ , so a natural candidate to consider for  $t = 8$  is  $(X, F) = [101100110011]$ . Setting accordingly  $(X, F) = (12, \{1, 3, 4, 7, 8, 11, 12\})$  indeed yielded an improved upper bound  $c_8 \leq 0.7002 \times 2^{1-\binom{8}{2}}$ , see Table 5.

**Proposition 1.** *We have  $c_6 \leq 0.7445 \times 2^{1-\binom{6}{2}}$ ,  $c_7 \leq 0.6869 \times 2^{1-\binom{7}{2}}$ , and  $c_8 \leq 0.7002 \times 2^{1-\binom{8}{2}}$ .*

$i$	1	2	3	4	5
$S_i(X, F)$	505	125,010	14,562,090	726,780,600	13,191,935,400
$S_i(X, \bar{F})$	518	135,726	17,463,606	1,028,265,840	26,106,252,480

**Table 3.** The values of  $S_i(X, F)$  and  $S_i(X, \bar{F})$  when  $(X, F) = (10, \{1, 3, 4, 7, 8\})$

$i$	1	2	3	4	5	6
$S_i(X, F)$	1002	490,050	113,148,090	11,590,147,800	506,500,533,000	14,677,396,549,200
$S_i(X, \bar{F})$	1045	556,842	146,860,362	17,896,958,640	950,437,303,200	21,359,851,904,160

**Table 4.** The values of  $S_i(X, F)$  and  $S_i(X, \bar{F})$  when  $(X, F) = (11, \{3, 4, 7, 8, 10, 11\})$

$i$	1	2	3	4	5	6	7
$S_i(X, F)$	2,027	2,030,562	986,934,042	223,874,343,000	21,997,023,741,000	868,195,804,568,400	23,207,044,770,478,800
$S_i(X, \bar{F})$	2,068	2,158,860	1,120,464,444	279,763,013,640	32,608,321,954,560	1,762,344,151,444,800	47,296,455,155,389,440

**Table 5.** The values of  $S_i(X, F)$  and  $S_i(X, \bar{F})$  when  $(X, F) = (12, \{1, 3, 4, 7, 8, 11, 12\})$

## 5 Conclusion and future work

We presented a computational framework for computing the ratio of monochromatic  $t$ -cliques and the number of all  $t$ -subsets for a specific Cayley graph determined by a pair  $(X, F)$ . The program allows for searching for counterexamples to a 1960 Erdős's conjecture on multiplicities of complete subgraphs. We described a significant speedup obtained by the so-called  $m$ -approach and considering inherent symmetries. As a result, we were able to improve the known upper bounds for  $t = 6, 7$  and 8.

The computational framework presented lends itself to straightforward parallelizations. A parallel version of our program will allow us to explore larger  $t$ 's and also to enlarge the search space for smaller values of  $t$ . The first task thus will be to search for a better pair  $(X, F)$  for  $t = 8$  to improve the upper bound for  $c_8$ , and to redo the searches for  $t = 4, 5, 6$  and 7 in larger search spaces.



## Acknowledgements

The authors would like to thank the referees for their comments that helped improve the exposition. This work was supported by the Natural Sciences and Engineering Research Council of Canada, and by the Canada Research Chair program, and made possible by the facilities of the Shared Hierarchical Academic Research Computing Network (<http://www.sharcnet.ca/>).

## References

1. Conlon, D.: On the Ramsey multiplicity of complete graphs. *Combinatorica* (to appear).
2. Erdős, P.: On the number of complete subgraphs contained in certain graphs. *Mathematical Institute of the Hungarian Academy of Sciences* 7 (1962) 459-464.
3. Erdős, P., Moon, J.W.: On subgraphs of the complete bipartite graph. *Canadian Mathematical Bulletin* 7 (1964) 35-39.
4. Franek, F.: A note on Erdős' conjecture on multiplicities of complete subgraphs – lower upper bound for cliques of size 6. *Combinatorica* 3-22 (2002) 451-454.
5. Franek, F., Rödl, V.: Ramsey problem on multiplicities of complete subgraphs in nearly quasirandom graphs. *Graphs and Combinatorics* 8 (1992) 299-308.
6. Franek, F., Rödl, V.: 2-Colorings of complete graphs with a small number of monochromatic  $K_4$  subgraphs. *Discrete Mathematics* 114 (1993) 199-203.
7. Giraud, G.: Sur le problème de Goodman pour les quadrangles et la majoration des nombres de Ramsey. *Journal of Combinatorial Theory, Series B* 27 (1979) 237-253.
8. Goodman, A.W.: On sets of acquaintances and strangers at any party. *American Mathematical Monthly* 66 (1959) 778-783.
9. Jagger, C., Šťovíček, P., Thomason, A.: Multiplicities of subgraphs. *Combinatorica* 16 (1996) 123-141
10. Ramsey multiplicities, <http://optlab.mcmaster.ca/~jessiel/ramsey.htm>.
11. Sidorenko, A.F.: Cycles in graphs and functional inequalities. *Mathematical Notes* 46 (1989) 877-882.
12. Thomason, A.: A disproof of a conjecture of Erdős in Ramsey theory. *Journal of the London Mathematical Society* 39-2 (1989) 246-255.
13. Thomason, A.: Graph products and monochromatic multiplicities. *Combinatorica* 17 (1997) 125-134.